Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith.

Attorney Docket No.: 023070-125630US - Sheet 1 of 18

Wnt-1 Ab Æ MCF-7 SKBR-3 SW480 SA0S-2 LP-9 A549 8 8

F/G. /

Cell death after anti-Wnt Ab incubation (%)

## Fraction of apoptotic cell death (%) after anti-Wnt antibody treatment

Cell Line	No Treatment	Control Ab	Anti-Wnt-1 Ab	Anti-Wnt-2 Ab
CCL-75 (normal lung)	0.9 ±0.3	3.8 ±2.3	5.8 ±2.1	2 ±1.1
H838 (lung cancer)	1.7 ±0.8	3.1 ±3.0	28.4 ±4.6	58.3 ±8.1
H460 (lung cancer)	0.6 ±0.3	12.7 ±7.0	83.8 ±5.6	90.7 ±6.5
A549 (lung cancer)	0.3 ±0.1	4.9 ±3.0	84.7 ±3.3	82.3 ±4.5
MCF-7 (breast cancer)	0.7 ±0.4	1.6 ±0.7	56.7 ±3.9	47.2 ±4.2
SKBR-3 (breast cancer)	0.4 ±0.1	5.8 ±1.3	67.9 ±6.1	56.4 ±4.8
SW-480 (colon cancer)	5.9 ±0.8	14.6 ±4.6	43.3 ±4.4	48.9 ±5.2
SAOS-2 (sarcoma)	1.1 ±0.4	4.2 ±1.6	20.0 ±3.2	16.5 ±1.0
LP-9 (normal mesothelial)	2.3 ±1.1	5.1 ±2.7	4.1.±1.5	6.2 ±2.6
REN (mesothelioma)	13.6 ±2.4	15.2 ±6.5	89.6 ±3.8	81.4 ±8.4
H28 (mesothelioma)	6.6 ±3.5	12.4 ±7.4	68.7 ±5.8	41.3 ±6.6

FIG. 2

Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 02307O-125630US - Sheet 3 of 18




FIG. 3A







FIG. 3B







Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 02307O-125630US - Sheet 6 of 18

## FIGURE 5






H460-control mAb (50 μg) H460-anti-Wnt-1 mAb (50 μg)




Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 02307O-125630US - Sheet 9 of 18



in vivo study on breast cancer. Tumor volume is shown after 3 weeks treatment with anti-Wnt-1 monoclonal antibody and control monoclonal antibody. 5 animals are in each group. None of the animals grows tumor after anti-Wnt-1 mAb injection, but three out of five animals grow tumor in the control group. (I.P. injection was done once weekly one week after inoculation of the MCF-7 cells)

FIG. 6C

Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 02307O-125630US - Sheet 10 of 18



in vivo study on breast cancer. Tumor volume is shown after 3 weeks treatment with anti-Wnt-1 monoclonal antibody and control monoclonal antibody. 5 animals are in each group. None of the animals grows tumor after anti-Wnt-1 mAb injection, but three out of five animals grow tumor in the control group. (I.P. injection was done once weekly one week after inoculation of the MCF-7 cells)

FIG. 6C

Figure 7

1) CDR and FR region (light chain kappa) amino acid sequence of a anti-WNT1 antibody (Seq ID No:4)

Clone #1	V			!	!	!	Ĺτι	ĸ	н	1	н	Σ	ტ	H	 	 	1	; ; ;	1			:	
1921	1 D GAC	I ATT	v GTG	r CTG	5 T ACA	CAG	S TCT	CCT	A GCT	10 S TCC	L	A GCT	V GTA	S TCT	15 L CTG (	9 9 9	O CAG 1	R AGG (	A Scc	20 T ACC P	I ATC	s TCA	
	>			, v	×	ď	>	30					3.5 S. 50	>			1	40 H	1	ı	1	o	
1y21	TAC	A.	325	AGC	4	AGT	<b>7</b> 1	_	ACA	TCT		TAT		TAT	:		ATG (	<i>t</i> )	TGĠ	AAC . 0	CAA (	CĀĞ	
	ĮT.	æ	7	ı	н	Σ	v	H	!	!	^ !				CDR2	İ	IMGT				· 	! ! V	
1921	45 K AAA	PCCA	g GGA	CAG	P CCA	50 P	R AGA	r CTC	CTC	I ATC	SS Y TAT	L	V GTA	S TCC	:		:		:	:	65	N AAC	
	i		1		1	1					!	-		Ĺτι	pz;	m	1	н	Σ	ღ	۲	!	
1y21	L	e gaa	S TCT	70 G GGG	V GTC	P	:	A Scc	75 R AGG '	F TTC 1	S AGT	စ ဗိမ္မ	SAGT	80 0 0 0 0	•	:	s TCT (	2 599 2 6	85 T ACA (	D GAC 1	F TTC /	TACC	
1y21	L J	90 N AAC	I	H	P	v GTG	95 E GAG	E GAG	E GAG	D GAT (	A GCT	100 A GCA	T ACC	Y TAT	Y TAC	TGT o	105 Q CAG (	H CAC 7	IATT	CDR3	- II E GAG (	IMGT 110 L CTT	

										•					
			1 1 1	ഗ	TCA		ø	CAG	-		N AAC	1 1 1		۲	ACC
			!	н	AŢC		ø	CAA		65	:	E		Œ	TIC
				20 T	ACC	1	z	AAC			:	ტ		Ω	GAC
			!	ø		1	3	TGG			:	Σ	85	) <del>[</del>	ACA
			!	œ	AGG	 	40 H	CAC		}	:	н		O	999
			!	œ	CAG		Σ	ATG	IMGT		:	ı		ഗ	TCT
				ບ	999			:	1	09	:	ო			:
			1	15 L	rn.			:	CDR2		:	æ			:
			H	ω	TCT		¥	TAT			S TCC	Ţ	0	8 0	999
				>	GTA		35 S	AGT			V GTA			ഗ	AGT
ი გ	ອ		Σ	Ø	GCT	IMGT	×	TAT			LCTT	1			ggc
z	AAC		н		TTA	1	Ö	၁၅၅	^	25	Y TAT			ഗ	AGT
120 K	AAA		•	10 S	TCC	CDR1	ഗ	TCT			I ATC			ᄕᅺ	TTC
* 8	TGA		н	Ø	GCT		F	ACA			CTC		75	ž ¤	AGG
လ	AGC		ĸ	Ωι	CCT		30	AGT	H		CTC			Ø	၁၁၅
а (	S CO		Ľτ	တ	TCT		>	GTC	ღ		r Aga				:
ဗ	GGA			O	CAG		တ	AGT	Σ	20	P CCC			Д	CCT
115 G	<b></b>			2 T	ACA		×	AAA	н		P CCA			>	GTC
E	GAG			>	GTG	^ !	့တ	PC,	1		CAG	i	6	2 0	999
တ	TCG		1	>	GTG		25 R A	gcc	8		GGA			ഗ	TCT
22	CGT			н	ATT		æ	AGG	œ		д Б			团	GAA
E &	ACA			l D	GAC	į	×	TAC	Ēų	45	A A A	į		ы	CTA
	1 <b>y</b> 21	clone #2			ly22W1lkRs GAC ATT			ly22W11kRs TAC			K ly22W11kRs AAA				ly22W11kRs CTA

	<u> </u>					
	, see					
90 N I H P V E E E AAC ATC CAT CCT GTG GAG GAG	115 R R G E P S * N K R CGG AGG GGG GAG CCA AGC TGA AAT AAA CGG	FR 1 - I M G TF	1 5 20 D I V L T Q S P A S L A V S L G Q R A T I S 3AC ATT GTG CTG CCT CCT TCC TTA GCT GTA TCT CTG GGG CAG AGG GCC ACC ATC TCA 	Y R A S K S V S T S G Y S Y M H W N Q Q TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT GGC TAT AGT TAT ATG CAC TGG AAC CAA CAG	F R 2 - I M G T	FR3 - IMGT
L ly22W11kRs CTC	S T X ly22WllkRs AGC ACG TTX	No:9)	1 D ly23w2lkRs GAC	ly23w2lkRs T	F 45 K Iy23w21kRs AAA	1

Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 023070-125630US - Sheet 14 of 18

TGC AGC CDR3 Jy23w2lkRs ACA TTA GGG AGC TTA CAC GTT ACG GAG GGG GGA CCA AGC TGA AAA AAC GG TGC CTG 80 Q GGA GGA GGA CCI CCT CAG ACA TCC ATG CCT TCA CAC CTT

a anti-WNT1 antibody (Seg ID No:4) 3) CDR and FR region (heavy chain IgG1) amino acid sequence of


GATTCACITITAATACCIACGCC......AIGAACIGGGICCGCCAGGCICCAGGAAAGGGITIGGAAIGGGI TGCTCGCATAAGAACTAGACGTTATAATTCTGCAACATATTATGCCGATTCTGTGAAA....GACAGGTTCACCATCTCC XGTTXCAGCCTGXAGGAGTCXGGTGGA....GGATTGGTGCAGCCTAAAGGGTCATTGAAACTCTCATGTGCAGCCTCTG AGAGATGATTCACGGGGCATGCTCTATCTGCAAATGAACAACTTGAAAACTGAGGACACAGCCATGTATTACTGTGTGA a anti-WNT2 antibody (Seq ID No:9) 4) CDR and FR region (heavy chain IgG1) amino acid sequence of

Ö Σ

Ø	TCC		CAG CAG			×	TAC	;	×	TAC	 110 C
Σ	ATG	1	K AAG			65	:	Н	Æ	೦೦೮	CDR3 D GAT
20 K	AAG		V GTG				:	Ö	E	ACA	999
>	GTG	1	₩ TGG			E	ACT	Σ	85 N	AAC	W TGG
ď	TCA		40 S AGC			ß	AGT	Н	တ	TCC	R AGA
	GCT		J T		IMGT	ტ	GGT	ı	တ	TCC	105 A GCA
ď	999			•	1	¥ ¥	TAT	m	×	AAA	C C TGT
15 P	CCT				CDR2	ט	GGA	œ	Ω	GAC	F
×	AAG			•		Д	CCT	ſτι	80 A	GCT	Y
٥	GTG		35	· ·		×	TAT		H	ACT	V GIC
۲	CTG	IMGT	٥ ليان	;		н	ATT		ы	CTG	100 A GCG
[±	GAG	1	7. Y	•	•	55 E	GAG		E-	ACA	S
10	:	CDR1	D 6			Ø	GGA		4	၁၁၅	D
۵	CCT		F F			н	ATT	į	75 X	AAG	GAG
C	GGA		30 FF			3	TGG	i	ď	399	S
;	JCX		ΕΗÇ	<b>5</b> 5		ы	GAG	į		:	95 T ACA
;	AG.		>	Σ		20 L	CTT		×	AAG	r CTG
rs.	:		ဗ ဦ	<b>5</b> H		Ö	GGC		(z.	TTC	S AGC
	:	^ !	s E	י ל		0	CAG		70 K	AAG	S
	:		25 A			ღ	GGA		[£	GAG	CTC
	:		¥ \$	8 8		H	М,	1	2	TAC AAT	90 O CAG
Н	:		ပင္ပ			45 ጸ	r.	1	>		MATG
	ly25W2HgRs		6. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	ıyzəmznyks			ly25W2HgRs			ly25W2HgRs	ly25W2HgRs

Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 023070-125630US - Sheet 16 of 18

FIGURE 8



Title: METHODS FOR TREATING CANCER BY INHIBITING WNT SIGNALING Inventors: Bia HE et al. – Filed: Herewith Attorney Docket No.: 023070-125630US - Sheet 17 of 18

FIGURE 9




Figure 10. Over-expression of Wnt signal antagonist (FRP or DKK) induces apoptosis in cancer cell line MS-1

